

## **A Systematic Review of the effect of Dietary and Nutritional Interventions on the Behaviours and Mental Health of Prisoners**

Matthew Poulter<sup>1</sup>, Shelly Coe<sup>1</sup>, Catherine Anna-Marie Graham<sup>1,2,3</sup>, Bethan Leach<sup>4</sup>, Jonathan Tammam<sup>1</sup>

<sup>1</sup>Centre for Nutrition and Health, Faculty of Health & Life Sciences, Oxford Brookes University, Oxford, United Kingdom, OX3 0BP.

<sup>2</sup>cereneo Foundation, Center for Interdisciplinary Research (CEFIR), 6354 Vitznau, Switzerland.

<sup>3</sup>Lake Lucerne Institute, 6354 Vitznau, Switzerland.

<sup>4</sup>Practice Plus Group, Hawker House 5-6 Napier Court, Napier Rd, Berkshire RG1 8BW

**Corresponding author:** M. Poulter, Email: matt0778@gmail.com, Centre for Nutrition and Health, Faculty of Health & Life Sciences, Headington Campus, Oxford Brookes University, Oxford, United Kingdom, OX3 0BP

**Short title:** Dietary Interventions on Mental Health and Behavioural outcomes in prisons.

**Ethical Standards Disclosure:** This work did not involve the use of human participants.



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114524000849

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

**Abstract**

Prisoners experience a higher burden of poor health, aggressive behaviours and worsening mental health than the general population. This systematic review aimed to identify research that used nutrition-based interventions in prisons, focusing on outcomes of mental health and behaviours. The systematic review was registered with PROSPERO on the 26th of January 2022: CRD42022293370. Inclusion criteria comprised of current prisoners with no limit on time, location, age, sex, or ethnicity. Only quantitative research in the English language was included. PubMed/Medline, Web of Science, EMBASE, PsycINFO, and CINAHL were searched, retrieving 933 results, with 11 included for qualitative synthesis. Studies were checked for quality using the ROB 2 or ROBINS-I tool. Of the included studies, seven used nutritional supplements, three included diet changes, and one used education. Of the seven supplement-based studies, six included rule violations as an outcome, and only three demonstrated significant improvements. One study included mental health as an outcome, however, results did not reach significance. Of the three diet change studies, two investigated cognitive function as an outcome, with both reaching significance. Anxiety was included in one diet change study, which found a significant improvement through consuming oily fish. One study using diet education did not find a significant improvement in overall mental resilience. Overall, results are mixed, with the included studies presenting several limitations and heterogeneity. Future research should aim to consider increased homogeneity in research design, allowing for a higher quality of evidence to assess the role nutrition can play in improving the health of prisoners.

## Introduction

Malnutrition can significantly affect one's ability to achieve optimal health and well-being<sup>(1)</sup>. Research has consistently identified the relationship between nutrition and health, demonstrating that inadequate nutrition could have detrimental effects at any stage of life<sup>(2)</sup>. From early foetal development, a pregnant mother's dietary choices can play a pivotal role in preventing undernutrition in her child and influence the risk of obesity in the child's adulthood<sup>(3)</sup>. Moving into adulthood, nutrition continues to play an important role in maintaining many aspects of health, where selected micronutrient deficiencies, such as a lack of vitamin D and calcium, could lead to an increased risk of osteoporosis<sup>(4)</sup>. These examples set a basis of evidence of deficiencies impacting health outcomes, though it is important to recognise that malnutrition does not solely refer to a lack of macro and micronutrients but also to overnutrition, such as a diet high in energy intake, fat, sugar, and salt<sup>(5)</sup>. These are key health issues in many developed countries, where access to food is generally plentiful and is often overprocessed with fat, sugar, and salt<sup>(5)</sup>. A diet high in these could increase the risk of obesity and obesity-related diseases such as diabetes, cancer, and cardiovascular disease<sup>(4, 6, 7)</sup>. While many of the identified effects of malnutrition relate to physical health and development, there is also emerging evidence of the impact of nutrition on mental health<sup>(8)</sup>.

Mental health is considered a state of mental well-being that enables people to cope with the daily stresses of life and contributes to their ability to work and engage with society and their community<sup>(9)</sup>. It is estimated that 970 million people worldwide live with at least one mental health disorder, with anxiety and depression being the most common<sup>(10)</sup>. Other prevalent mental health disorders include bipolar disorder, post-traumatic stress disorder, schizophrenia, eating disorders, and disruptive behavioural disorders<sup>(10)</sup>. Health organisations and associations, including the NHS<sup>(11)</sup>, the Association of UK Dietitians<sup>(12)</sup>, and the Royal College of Psychiatrists<sup>(13)</sup>, recommend a healthy diet as a crucial aspect of living with mental health disorders. This recommendation aims to aid recovery and ensure overall health while living with a disorder. Several reviews have identified the effects of diet on mental health outcomes, suggesting limited evidence of a relationship.

One such review identified that a Mediterranean diet, characterised by a high intake of fruits, vegetables, whole grains, and fish, with low consumption of meat and dairy, was inversely associated with the risk of depression<sup>(14)</sup>. However, this review noted caution with the results due to various methodological differences between the included studies. A later meta-analysis

further identified the positive effects of a healthier diet on the risk of depression<sup>(15)</sup>. In addition, an unhealthier Western-style diet, characterised by a high intake of red meat, sweets, and processed foods, is associated with a higher risk of depression. However, similar to the previous findings, there was heterogeneity among the studies included, and the authors stated that further research is needed to confirm these findings<sup>(15)</sup>. Nonetheless, these reviews provide a basis of evidence for a relationship between mental health and diet and the growing interest in this field. Yet, despite the body of evidence supporting nutrition's role in human health, development, and behaviour, select population groups remain vulnerable to malnutrition.

Vulnerable populations are defined as those who have limited resources to mitigate potential challenges in their lives, including challenges to their health outcomes<sup>(16)</sup>. Examples of vulnerable populations can include racial minorities, those living in poverty, the elderly, the homeless, people with mental health disorders, and those in contact with the justice system, such as prisoners<sup>(16, 17)</sup>.

Globally, the prison population is estimated to be approximately 10.8 million, though this figure could be as high as 11.5 million when accounting for incomplete data, as reported by the World Prison Population List<sup>(18)</sup>. Prisoners are considered a vulnerable population, often experiencing high levels of poor mental health and aggressive behaviours attributable to numerous factors. Examples include a life of poverty, past trauma, low educational achievements, and poor general health<sup>(19-21)</sup>. The current population of UK prisoners stands at 88,126 as of the 20th of October 2023, with their mental health reported as highly compromised<sup>(22)</sup>. It has previously been reported that only around 10% of prisoners receive treatment for mental disorders, while up to 70% are estimated to suffer from at least one undiagnosed disorder<sup>(20)</sup>. The prison environment can be detrimental to health due to overcrowding, lack of purposeful activities (i.e., work/education), and inadequate nutritional intake<sup>(19, 23-25)</sup>.

The World Health Organisation (WHO) sets out that prisoners are entitled to basic human rights, including access to nutrition. However, evidence identifies that prisoners are either choosing a diet that fails to meet reference nutrient intakes, or prison institutions are not providing adequate options to meet nutritional recommendations. This issue has been identified in multiple countries<sup>(19, 25-29)</sup>. We recognise that nutrition is not necessarily a

solution to solve all health problems facing prisoners; however, good nutrition forms a key part of overall health and well-being. Research must aim at addressing nutritional needs, which have the potential to improve overall prisoner health, including mental health outcomes.

Therefore, this review aims to systematically examine dietary and nutritional interventions implemented in prisons among the prisoner populations to investigate their impact on mental health and a spectrum of behaviours. The term 'behaviours' is intentionally kept broad, encompassing a range of actions that may influence the overall well-being of prisoners. Notably, this includes behaviours linked to mental health, such as self-harm and aggression<sup>(30-32)</sup>. While presenting one of these behaviours alone is not enough to necessarily identify a mental disorder, in some cases, it can indicate an underlying undiagnosed mental disorder. Furthermore, disruptive behaviours are considered under the umbrella of mental health disorders, and the WHO notes that risk-taking behaviours such as substance abuse and violence can indicate underlying emotional issues<sup>(10, 33)</sup>. This comprehensive approach in the review acknowledges the interconnectedness of mental health and behaviours, recognising both the prevalence of undiagnosed mental health concerns and the escalating levels of aggressive and violent behaviours observed across prisons globally<sup>(34-39)</sup>.

## **Methods**

This systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA) 2020<sup>(40)</sup>. This review was registered in the International Prospective Register of Systematic Reviews (PROSPERO; registration ID: CRD42022293370), where the study protocol and amendments can be found.

### *Search Strategy*

Five electronic databases were searched: PubMed/Medline, Web of Science, EMBASE, PsycINFO, and CINAHL, with no restriction on the publication date, using the filter for the English language. The initial search was conducted on the 3rd of March 2022, with follow-up searches on the 7th of June 2022 and the 18th of October 2023 to ensure this review captured any new research published during the writing of this paper. The search strategy included key terms related to prison/prisoners, mental health, behaviours, and diet interventions. Key terms were combined with Boolean operators 'AND' and 'OR'. See Appendix 1 for full search terms. All retrieved results were organised in Microsoft Excel.

### *Inclusion and Exclusion Criteria*

Inclusion: (1) peer-reviewed articles published in English during any year globally; (2) study designs included are randomised controlled trials (RCTs), cohort studies, and case-control studies; (3) the study population must be prisoners at the time of the study, no restrictions on age, sex/gender, ethnicity or sexuality were applied; (4) any diet/nutrition based intervention was considered; (5) any study where the outcome examined mental health or behaviours were included; (6) studies must include a control or comparator group. Exclusion: (1) studies which have not been peer-reviewed, and which are not in the English language; (2) qualitative and mixed method study designs; (3) populations which were not prisoners at any point during the study period; (4) non-diet-based interventions; (5) physical health outcomes; (6) studies lacking control or comparator.

### *Data Extraction*

Data extraction began on the 1st of September 2022, with one reviewer extracting data, a second reviewer checking the work, and any disagreements discussed. A third reviewer was consulted if necessary. Key data extracted included: (1) general study characteristics: author(s), year, intervention type, location, group studied, and study length; (2) characteristics of participants: number of participants, sex, age, and inclusion/exclusion criteria; (3) intervention details: exposure, control/comparator, outcome domain, outcome measurement tools; and (4) results: key study results, including any effect measures (e.g., mean difference, odds ratio), and overall conclusion.

### *Quality Assessment*

Two reviewers independently evaluated the quality and risk of bias for the included studies. Risk of bias and quality assessment for randomised studies was conducted using the revised tool to assess the risk of bias in randomised trials (RoB 2) tool<sup>(41)</sup>. The RoB 2 tool assesses bias based on five domains: the randomisation process, deviations from the intended intervention, missing outcome data, outcome measurement, and selection of the reported result.

The risk of bias in non-randomised studies of interventions (ROBINS-I) tool was used for non-randomised studies<sup>(42)</sup>. The ROBINS-I tool measures bias based on seven domains: confounding, participant selection, intervention classification, deviations from intended intervention, outcome measures, and results reported.

### *Data Synthesis*

A meta-analysis was considered for this review; however, due to the anticipation of a diverse range of study types and methodologies, it was deemed to be of no benefit. Qualitative synthesis was conducted, where studies were grouped based on their intervention type and outcome. Additionally, due to the terms 'prisoners/inmates/offenders' being used interchangeably, for this review, only the term 'prisoner/s' will be used.

## **Results**

### *Study Selection*

The database search identified 933 results, leaving 714 for title screening after duplicates were removed. After title screening, 633 records were removed with a further 60 removed following abstract screening. An additional paper was identified through hand-searching literature. The follow-up search on the 18th of October 2023 resulted in no further papers identified. In total 22 results for full-text review were identified, after which 11 were deemed eligible for inclusion in qualitative synthesis, see Figure 1 for the PRISMA flow diagram<sup>(40)</sup>. See Table 1 for included study characteristics. See Appendix 2 for the excluded studies' general characteristics and reasons for exclusion.

## **Results**

### *Quality Assessment*

Of the 11 included studies, nine were assessed using the ROB 2 Tool. Of these, eight were found to have low overall bias, while one was found to have some concerns (see Appendix 3). Two studies were assessed using the ROBINS-I Tool, and both were found to have critical bias (see Appendix 4).

### *General Study Characteristics*

Of the eleven included studies, seven were conducted in the USA<sup>(43, 45, 47-49, 51, 52)</sup>. One study came from each of The Netherlands<sup>(53)</sup>, Singapore<sup>(50)</sup>, United Kingdom<sup>(46)</sup>, and Australia<sup>(44)</sup>. The earliest published study was from 1975<sup>(45)</sup>, with the next being in 1994<sup>(43)</sup>, and the remaining studies were all published after 2000. Study length varied, ranging from 3 days<sup>(43)</sup> to 12 months<sup>(50)</sup>. Of the eleven studies, nine were RCTs<sup>(43, 44, 46-48, 50, 52, 53)</sup> with the remaining two being non-randomised<sup>(45, 49)</sup>.

### *Participant Characteristics*

The number of participants in a study varied between 25 and 449. The supplementation intervention studies included the highest number of participants<sup>(44-46, 50-53)</sup>. Only two studies included female participants<sup>(49, 51)</sup>, one of which included both male and female participants<sup>(51)</sup>. The age of participants was not reported in two studies<sup>(45, 46)</sup>, with one study only reporting the age range and not the mean age of participants<sup>(43)</sup>.

### *Interventions, Outcomes, and Results*

The majority of studies, seven, were supplementation-based<sup>(44-46, 50-53)</sup>, with only three assessing the impact of diet change<sup>(43, 47, 48)</sup>, and one investigating diet education<sup>(49)</sup>. See Table 2 for a summary of exposures, controls/comparators, outcomes, and key results for the included studies.

### **Effect of Supplements on Behaviours**

Six supplement-based studies investigated behavioural outcomes, including rule violations and aggression. Four studies demonstrated a significant reduction in rule violations<sup>(46, 51-53)</sup>, and only one study found a significant reduction in aggression<sup>(50)</sup>. Information on the types of supplements used in each study can be found in Table 2, and study characteristics in Table 1. In a randomised placebo-controlled trial Gesch *et al*<sup>(46)</sup> found that the active supplementation group experienced a 26.3% reduction in the number of infringements compared to the placebo group ( $p=0.03$ ). A similar RCT by Zaalberg *et al*<sup>(53)</sup> conducted in the Netherlands also found that the number of recorded incidents of rule-breaking post-intervention was significantly less in the supplement group (incident rate ratio (IRR) = 0.6;  $p=0.017$ ). Schoenthaler *et al*<sup>(51)</sup> found in their RCT that for non-violent rule violations, there was an 83% decrease in the experimental group compared to 49% in the placebo group ( $p=0.008$ ). Violent rule violations fell 80% in the experimental group and 56% in the placebo group ( $p=0.44$ )<sup>(51)</sup>. Overall, total rule violations fell 83% for the supplement group and 55% in the placebo group (95% CI: 15-41%,  $p=0.005$ ). A more recent RCT by Schoenthaler *et al*<sup>(52)</sup> used high and low-dose multivitamin supplements, which found a statistically significant difference between the placebo and low-dose supplement group for serious rule violations (RR=0.61, 95% CI: 41-90,  $p=0.01$ ). The result between the placebo and high-dose group was not statistically significant (RR=1.48, 95% CI:0.99-2.23,  $p=0.06$ )<sup>(52)</sup>. Two of the more recent studies found no significant results for rule violations<sup>(44, 50)</sup>.

Three of the supplementation studies investigated aggressive/antisocial behaviours<sup>(44),(50),(53)</sup>. Raine *et al*<sup>(50)</sup> conducted an RCT resulting in the omega-3 supplement group showing significantly lower self-reported antisocial behaviours compared with the placebo group after



the 3-month experimental period ( $p=0.006$ ;  $d=0.56$ ). At 3 months post-intervention, the omega-3 supplement group continued to report lower levels of antisocial behaviour compared to the placebo group, though the result failed to reach significance ( $p=0.059$ ;  $d=0.39$ )<sup>(50)</sup>. At 6 months post-intervention, however, the omega-3 group reported significantly lower levels of antisocial behaviours compared with the placebo group ( $p=0.045$ ;  $d=0.43$ )<sup>(50)</sup>. No difference was found between groups based on officer reports of institutional infractions ( $p=0.36$ )<sup>(50)</sup>. Cortie *et al*<sup>(44)</sup> in a feasibility pilot RCT found that aggressive/antisocial behaviours were not significantly different through supplementation when assessing behaviours using the Inmate Behaviour Observation Scale (IBOS) ( $X^2(1) = 0.83$ ,  $p=0.36$ ), the self-report aggression questionnaire (AQ;  $X^2(1) = 0.03$ ,  $p=0.82$ ), and Brown's Attention Deficit Disorder Scale (BADDS;  $X^2(1) = 0.06$ ,  $p=0.64$ ). A subset analysis was also conducted with participants who were characterised as aggressive based on their IBOS baseline scores. Only 11 participants from the supplement group and 18 from the placebo were eligible for this subset analysis. In the supplement group, there was an improvement of 7% for the IBOS ( $p=0.566$ ), 25% for records of misconduct ( $p=0.103$ ), 19% for AQ ( $p=0.371$ ), and 2% for BADDS ( $p=0.910$ ), however, all failed to reach significance. Zaalberg *et al*<sup>(53)</sup> found that aggression, as measured by the staff-reported SDAS, demonstrated no significant difference between the supplement and placebo group ( $p=0.23$ ).

### **Effect of Supplements on Mental Health**

Two of the supplementation studies included outcomes around mental health, with mixed results<sup>(45, 53)</sup>. D'Asaro *et al*<sup>(45)</sup> conducted a non-randomised controlled trial, finding that supplements did decrease feelings of anxiety, however, the significance was not reported. Zaalberg *et al*<sup>(53)</sup> found, through the General Health Questionnaire-28 (GHQ-28), that there was a general trend for the supplementation group to experience better well-being; however, this did not reach significance ( $p=0.069$ ).

### **Effect of Diet Change on Mental Health**

Only one study which used diet change investigated mental health outcomes, where measures of anxiety improved<sup>(47)</sup>. Hansen *et al*<sup>(47)</sup> aimed to identify the effects of diet change on mental health using an RCT. Heart rate (HR) and heart rate variability (HRV), measured as the root mean square of successive differences (rMSSD), were used as biological indicators of anxiety. The State-Trait Anxiety Inventory (STAI) was used for self-reported anxiety. A diet change for the experimental group, which involved the provision of salmon three times per week for 23 weeks, resulted in a reduced HR for the experimental group from baseline to post-intervention ( $p<0.01$ ;  $d=0.45$ ). rMSSD increased in the experimental group from

baseline to post-intervention, indicating anxiety improved ( $p < 0.01$ ;  $d = 0.45$ ). Additionally, vitamin D status was significantly related to rMSSD post-intervention ( $p < 0.05$ ;  $r = 0.27$ ), though not at baseline ( $p > 0.05$ ). The STAI revealed there was a significant decrease in state anxiety ( $p < 0.03$ ;  $d = 0.43$ ) for the experimental group, but no significant result was identified for trait anxiety ( $p > 0.05$ ).

### **Effect of Diet Change on Cognitive Function**

One study investigated the effects of diet change on cognitive function, which found that diet change improved cognitive function<sup>(48)</sup>. Hansen *et al*<sup>(48)</sup> conducted an RCT, **that** was a part of the same diet change study by Hansen *et al*<sup>(47)</sup>, which involved the provision of fatty fish three times a week to the experimental group. Cognitive function was measured by the Tower of Hanoi task (ToH), measuring executive function (i.e., the ability to plan and solve tasks), and the Iowa Gambling Task (IGT), measuring risky decision-making. For the ToH task, the experimental group improved from baseline to post-intervention ( $p = 0.016$ ;  $d = 0.35$ ). A subgroup analysis was conducted with participants who had a previous history of drug/alcohol abuse, though no significant improvements in either group for the IGT were found ( $p = 0.109$ ;  $d = 0.41$ ). The subgroup analysis for the ToH showed that only those in the experimental group with a history of previous drug/alcohol abuse improved from baseline to post-intervention ( $p = 0.013$ ;  $d = 0.45$ ). Correlations between the nutritional status of EPA, DHA, and vitamin D and cognitive performance were also conducted. No correlation between vitamin D status and cognitive performance was found. The effect of the sum of EPA and DHA on the sub-task with low working memory load task was found at baseline ( $p = 0.001$ ;  $r = 0.41$ ), though not at post-intervention ( $p = 0.858$ ;  $r = 0.02$ ).

### **Effect of Diet Change on Behaviour**

One study investigated the effects of diet change on various behavioural outcomes including aggression, and impulsivity, resulting in improved behaviours<sup>(43)</sup>. Bachorowski *et al*<sup>(43)</sup> used diet change to identify the effects of sucrose on neuropsychological test performance, in three groups of juvenile offenders based on their serum glucose nadirs (low, borderline, and normal). Participants ingested a sucrose-loaded breakfast, and a no-sucrose control breakfast before a battery of neuropsychological tests, in a double-blind randomised crossover design. Participants were also observed by the researchers, who used a 32-item behaviour checklist to evaluate observations of participants regarding their attention and motivation. The checklist was completed by the researchers after each day of neuropsychological tests and was formed from six subscales looking at restlessness, distraction, aggression, impulsivity, awkwardness, and insecurity. It was found that there was an interaction between **the** nadir serum glucose

group, order of breakfast and subscale of the behaviour checklist ( $p < 0.05$ ). The results indicated that participants with low and normal nadirs were rated as having better behaviour following the sucrose-loaded meal, though overall the study suggests the relationship between nadir serum glucose levels, breakfast type, and behaviour improvement is complex.

### **Effect of Diet Education on Mental Health**

One study used diet education as the intervention, which investigated whether the mental resilience of participants would increase following the education course, however, no significant improvements were found<sup>(49)</sup>. Johnson et al<sup>(49)</sup> conducted health education classes, comprising three sessions over three months each lasting between 60-90 minutes, as part of a quasi-experimental pilot, which focused on dietary education i.e., making healthier food choices, portion control etc. The results showed a slight improvement in mental resilience between baseline, six weeks, and 12 weeks. However, these improvements did not reach significance ( $p = 0.437$ ).

## **Discussion**

### **Overview of the Results**

The aim of this review was to identify existing literature, which reports the effects of nutritional interventions on the mental health and behaviours of prisoners. A previous systematic review was identified, however, this only looked at the effects of supplements on aggression outcomes<sup>(54)</sup>. Our review differed by considering all types of nutritional interventions on all mental health and behaviour outcomes. From the 11 studies included three types of nutritional interventions appeared: diet change, diet education, and dietary supplements.

### **Dietary Supplements**

Supplementation studies were the most prevalent type of intervention in prisons, likely due to the ease of administering the intervention to a large population, in contrast to the more logistically and time-intensive methods of education and diet change. Although supplements were found to have significant positive results, there was a degree of heterogeneity across the studies' overall conclusions.

Regarding the types of supplements used, three studies employed a combination of vitamin-mineral and fatty acid supplements<sup>(44, 46, 53)</sup>. Raine et al<sup>(50)</sup> focused solely on the use of a fatty acid omega-3 supplement. Two studies, D'Asaro, Groesbeck, and Nigro<sup>(45)</sup> and Schoenthaler et al<sup>(51)</sup> utilised only a vitamin-mineral supplement. Schoenthaler et al<sup>(52)</sup> offered either a low-dose vitamin-mineral supplement meeting 100% of the Recommended Dietary Allowance (RDA) or a high-dose supplement exceeding the RDA for certain vitamins. D'Asaro,

Groesbeck, and Nigro<sup>(45)</sup> provided only vitamin-mineral supplements to correct existing deficiencies within the sample population. Refer to Table 2 for a summary of the interventions used in each study, including supplement details.

### ***Dietary Supplements, Rule Violations, and Aggression***

Rule violations emerged as the predominant outcome measured, with six out of the seven supplement studies incorporating this parameter. This preference is likely attributed to the accessibility of such data, as prison staff are generally mandated to report all rule violations, encompassing both violent and non-violent incidents, such as prisoner refusal to work. The evidence that shows the impact of supplements in reducing rule violations is largely positive. However, it's noteworthy that four studies reported significant effects with similar medium effect sizes, emphasizing the need for larger sample sizes to substantiate these findings<sup>(46, 51-53)</sup>.

A key issue with the supplementation studies was identifying why any improvement in the outcomes measured may have occurred. For example, why rule violations may have reduced over the course of supplementation. This uncertainty is largely due to the variability between the supplements used in the studies. For example, Zaalberg *et al*<sup>(53)</sup> replicated the study by Gesch *et al*<sup>(46)</sup>, yet the quantities of vitamins, minerals, and fatty acids provided differed significantly in many cases. Both studies investigated omega-3 fatty acids, yet Zaalberg *et al*<sup>(53)</sup> provided almost five times as much omega-3 fatty acids as Gesch *et al*<sup>(46)</sup>, and half the amount of vitamin D. As previously discussed, the evidence implicating omega fatty acids is limited when regarding mental health outcomes. However, given that rule violations can be linked more towards feelings of aggression and impulsivity, there is a potential that the underlying effects of omega deficiencies could explain the improvement seen. Previous research has identified some implications towards omega fatty acids regarding this, both in prison populations and children<sup>(55, 56)</sup>. The two studies<sup>(44, 50)</sup> included in this review which identified no significant reductions from omega fatty acid supplements may have additional confounders leading to non-significant results.

The first of these two studies by Raine *et al*<sup>(50)</sup> conducted their study in Singapore, with participants mostly being of Chinese or Malaysian descent. Non-significant findings could relate to the fact that Southeast Asian cuisine tends to contain higher levels of fish, and in turn, omega fatty acids when compared to Western diets<sup>(50, 57)</sup>. Therefore, at baseline, this population would be less likely to have omega-3 deficiencies, and supplementation would offer little benefit. However, there is no baseline dietary intake data provided, a common theme in the included literature. Therefore, it cannot be stated with certainty that the

participants were omega-3 sufficient at baseline. Furthermore, cultural factors are reported, where Singaporean institutions report fewer serious rule violations, particularly relating to drugs and alcohol, in comparison to studies in the Western world<sup>31</sup>. Therefore, it is conceivable that a less violent population would be less likely to demonstrate the potential effects of omega-3 in reducing violent or antisocial behaviours. Though a decrease in rule violations was not observed, Raine *et al*<sup>(50)</sup> demonstrated that self-reported aggression was reduced. Similarly, Cortie *et al*<sup>(44)</sup> did not find results that meet significance, for rule violations or self-report aggression, though there is a trend for rule violations to decrease in the supplement group. However, it is not possible to fully test the accuracy of this result due to a small sample size (n=131), with a post-hoc power calculation finding that a sample size of n=600 would be needed to detect any significant effects.

Two studies that observed a reduction in rule violations used only vitamin-mineral supplements<sup>(51, 52)</sup>. This leads us to consider whether the effects seen across the supplementation studies were attributed to a mechanism involving the vitamin and minerals component. Vitamin D is notorious for being deficient in populations globally, and in the context of a prison, this deficiency is more likely due to limited access to sunlight<sup>(27, 58, 59)</sup>. Among the included studies, Gesch *et al*<sup>(46)</sup> found that vitamin D was deficient based on the baseline dietary assessment, with prisoners, on average, having an intake of 3.5µg, while the current dietary recommendations suggest 10µg per day<sup>(60)</sup>. Limited evidence suggests vitamin D's role in improving behaviours from studies involving animals and children, though further research is necessary<sup>(61, 62)</sup>. It is interesting to note that all the supplementation studies discussed in this review, included some amount of vitamin D. The effects of supplements identified in this review could be attributed to any number of vitamins, minerals, and fatty acids. This further highlights the common theme that the studies in this review lacked baseline dietary assessment, though this serves as an avenue for future research.

### ***Dietary Supplements and Mental Health***

Mental health outcomes were not a prominent feature in the included studies. Only one study included general well-being as an outcome measure, where despite a general trend of improvement, this ultimately did not reach significance<sup>(53)</sup>. However, this study suffered from a high dropout rate, reducing the sample size and potentially limiting the significance of the result.

The earliest study included in this review, from 1975, found a significant decrease in self-reported anxiety; however, no significant value was provided<sup>(45)</sup>. Due to the lack of data and unclear methodology reported, this study was challenging to interpret and, thus, overall was

found to be of high bias. High bias was in part due to the authors not reporting whether the outcome measures were suitable for the population. This study also had the smallest number of participants among the supplementation studies in this review, with a particularly small control group (n=7) compared to the experimental group (n=21), an issue pointed out by the authors. Participants in this study reported several side effects, including excessive hunger and skin eruptions. It is unknown if these were due to supplements or other reasons; however, no other study included in this review reported similar issues. Bias in the study was also discussed by the authors, noting that all participants were in the education program which detailed the study and its purpose, potentially creating a placebo effect and subject bias<sup>(45)</sup>.

Despite the high burden of mental health disorders within prisons, there was very little consideration of mental health outcomes in the included studies. Previous research outside of the prison population identified beneficial effects of supplementation on mental health outcomes, particularly for vitamin D and depression, and recent guidelines recommend the use of omega-3 for major depressive disorder<sup>(63-66)</sup>. Based on the limited evidence identified in this review regarding supplements and mental health, we conclude that research in this area is warranted.

### **Diet Change**

Two diet change studies resulted from a larger study on mental health and nutrition<sup>(47, 48)</sup>. Salmon was used primarily as a natural source of omega-3 fatty acids and vitamin D, considered more beneficial than supplement use by the authors<sup>(48)</sup>. Both of these studies provided significant positive associations with fatty fish consumption, where the experimental group experienced reduced biological indicators of anxiety, state anxiety, reduced impulsivity, risky decision-making, and improved planning and solving skills. Overall, results need to be interpreted with caution due to the nature of this type of dietary intervention, as issues lie in the inability to blind subjects. The authors reported this as a limitation and stated that both groups could have been influenced by the meals served, potentially perceiving salmon or chicken control as the healthier option.

Previous research outside the prison population has been unable to demonstrate a significant effect of fatty fish intake on rMSSD and HRV<sup>(67)</sup>. Furthermore, a meta-analysis examining the effects of omega-3 and polyunsaturated fats on anxiety and depression symptoms found little evidence to suggest a significant effect for preventing or reducing depressive or anxiety symptoms<sup>(68)</sup>. This meta-analysis highlighted a major limitation of the studies included, as many lacked a baseline dietary intake analysis – a limitation shared by the two diet change studies in this review. However, blood samples to monitor omega-3 blood levels were taken



to identify whether these had improved post-intervention for the experimental group. A more recent paper updating clinical guidelines around treating psychiatric disorders with nutritional therapies did recommend the adjunctive use of omega-3 for those with major depressive disorder<sup>(63)</sup>. This recommendation is based on the results of five RCTs and a statistically significant meta-analysis. Without understanding the overall baseline dietary status of participants, it is challenging to deduce what may be causing the significant effects, given that salmon provides more than omega-3, including vitamin D, selenium, and iodine, and is a source of protein. Whether the underlying biological mechanism was down to one or more of these other nutrients is unclear, though further investigation is warranted. Vitamin D was identified as a potential cause for improvements, as the fatty fish used in this study provided around 5µg vitamin D, half the dietary recommendation for vitamin D. Positive associations were found between vitamin D status and rMSSD, indicating a potential role for vitamin D in improving anxiety. However, the authors reported that neither the control nor the supplement group were vitamin D deficient at baseline, requiring further investigation<sup>(47, 48, 69, 70)</sup>.

The third diet change study included in this review differed from the two previous studies discussed, as Bachorowski *et al*<sup>(43)</sup> aimed to identify the effects of sucrose on juvenile prisoners. This study did include an outcome focusing on behaviours; however, it was secondary to other outcomes which focused on neuropsychological tests not relevant to this review. Therefore, only the results of the behavioural checklist will be discussed. The authors had previously identified evidence that suggested an association between sugar consumption and behaviour, though the results were mixed<sup>(71, 72)</sup>. Additionally, a meta-analysis on this topic identified that overall there was little evidence supporting the implication of sugar on behaviour<sup>(73)</sup>. Bachorowski *et al*<sup>(43)</sup> demonstrated a small significant effect of sucrose on behaviour, though these results should be accepted with caution. This diet change does not reflect real-life sugar consumption and focuses on brief short-term ingestion of sucrose in a controlled environment. In addition, this study had a small sample size and was found to have concerns in the RoB-2. The results highlight that short-term sucrose ingestion produces beneficial results in juvenile prisoners. These include a reduction in distractedness and impulsivity seen across all three groups. Where the population of this study was prisoners, this could be seen as a desirable outcome. However, this study only considered short-term effects and future work would benefit from identifying the effects of sugar consumption over a longer period. More recent research in this area has further highlighted the short-term benefits of sugar consumption for juveniles, believed to be down to the mechanism of the juvenile brain requiring more glucose than an adult<sup>(74, 75)</sup>.

## Dietary Education

While the only study to use nutrition education did not yield significant results, it did find a general increase in mental resilience<sup>(49)</sup>. The researchers noted that many participants at baseline were already reported to have high levels of resilience, potentially limiting the degree to which an individual could experience an increase. Overall, this study was found to have a high bias due to the lack of control for confounders, the absence of a valid measurement tool, and the lack of a control group, which would aid in the validity of any results gained<sup>(76)</sup>.

Previous research on using education to improve mental health outcomes has yielded positive results, although it is challenging to make direct comparisons with this study due to differences in the measured mental health outcomes. For instance, a study focusing on nutrition education for obese women (n=44) in Japan, emphasizing gut microbiota education, significantly improved self-measured levels of depressive symptoms<sup>(77)</sup>. Another study implemented education among a population of children in China (n=171), aiming to enhance self-measures of anxiety, and reported significant improvements<sup>(78)</sup>. These previous studies, however, featured larger sample sizes than the study by Johnson *et al*, as well as broader exclusion criteria to reduce the risk of confounding. Nevertheless, these studies demonstrate that education can be a powerful tool for improving mental health outcomes.

The author acknowledged this study as a pilot study and the first to examine resilience in a prison population. Therefore, there are numerous opportunities for future studies to build upon, including the utilisation of high resilience levels and current mental health treatment as exclusion criteria, employing a larger population size, and incorporating a control group for a more comprehensive before-and-after comparison<sup>(79, 80)</sup>.

## Limitations

The research studies included in this review have several limitations that impact the overall findings. It was commonly found that there was no baseline dietary intake analysis conducted. Four studies did conduct baseline dietary analysis, such as a food diary analysis<sup>(46)</sup>, omega-3 blood analysis<sup>(44)</sup>, food waste and blood analysis (vitamin A, E, C, B5, B12, thiamine, riboflavin, niacin, pyridoxine, folates, calcium, magnesium, copper, iron, zinc, selenium, manganese, and chromium)<sup>(51)</sup>, and blood analysis (vitamin A, B5, B6, E, thiamine, riboflavin, niacin, folic acid, iron, chromium, calcium, manganese, copper, selenium, zinc, and magnesium)<sup>(52)</sup>. Identifying underlying nutritional deficiencies in each study population could provide a clearer picture of which nutritional issues may be causing poor outcomes.



Despite including various study types and locations, the lack of homogeneity among the included studies limited the ability to directly compare the effects of interventions on outcomes through a meta-analysis. A sub-group metanalysis was considered with the supplementation studies, as there were seven included in this review. Ultimately, this was decided against due to the heterogeneity between supplementation studies. It was difficult to identify the precise effects of supplements due to the variety of supplements, and the combination of supplement types used in the research included in this review. It is impossible to ascertain whether any improvements were due to any specific vitamin deficiencies being corrected, or if the offering of fatty acids i.e., omega-3 were the cause, or if there was a synergistic effect of combined supplements.

Additionally, this review highlights the lack of females in research regarding prisoner populations.

Only one study focused solely on female participants<sup>(49)</sup>, with one other study including both females and males<sup>(51)</sup>. This reduces the generalisability of results across sexes and therefore future studies should consider the inclusion of females, particularly due to females having different nutritional needs to males<sup>(81)</sup>. Similarly, the majority of included studies recruited adult populations, with only four of the eleven studies included having recruited young offenders. As with females the nutrition requirements differ across the life course, and this is something which should be considered.

### **Future Directions**

The heterogeneity presented in the included studies, particularly amongst those using supplementation, creates difficulties in fully interpreting the results identified. A key theme which has emerged over the discourse of the literature reviewed is a lack of baseline dietary assessment. Future studies would benefit from conducting dietary analysis at baseline, to identify specific population group nutritional needs. This would help elucidate changes in participant nutrition, and in turn enable research to identify the underlying mechanisms behind potential beneficial results.

Factors which have also limited the effects of interventions include the lack of stricter study criteria, which was noted in particular for aggression outcomes. Recruiting populations which exhibit high levels of aggressive behaviour at baseline would enable researchers to observe a greater change, compared to populations with low aggression at baseline. Similarly, the results of Johnson *et al*<sup>(49)</sup> were impacted by a study population with high resilience, reducing the ability to identify improvements.

We would also encourage further research including female participants, as females were generally lacking in the included studies. The underrepresentation of females in the research is concerning, particularly within the context of health and nutrition as females require different nutritional needs to males<sup>(81)</sup>. While we recognise males make up the majority of prisoners globally, and therefore it is somewhat understandable, females do still account for 6.9%<sup>(82)</sup>. Additionally, there are various subgroups within prisons which were mostly absent from the included studies in this review, for example, pregnant women, those with chronic diseases, elderly prisoners, and different ethnic groups.

Finally, as this was the first review of its kind, and due to the heterogeneity of the included studies, both overall and within the three intervention groups identified, a meta-analysis at this stage was felt to be of little benefit. However, a future review would benefit by narrowing in on one of the intervention types identified, or conducting sub-group metanalysis for each intervention type once more emerging evidence is identified and published. This could allow for a meta-analysis, to provide improved estimates of the effects seen, and increase generalisability of study results.

### **Conclusion**

Nutrition is fundamental to a person's overall health and well-being, and nutritional interventions have been identified as a possible means to improve the prison environment, with the greatest effect seen in the reduction in the number of rule violations. However, this review identifies that the literature concerning prisoners, mental health, and nutrition is still in its infancy and that the results discussed should be interpreted with caution. We encourage further research in this area, to fully understand the effects nutrition can have, and this review sets out future directions which could be taken, to contribute to the benefit of the prison population.

**Acknowledgements:** The authors would like to acknowledge the work and collaborative efforts of Professor John Stein (University of Oxford), Alberto Crisci (HMPPS), Rupert Bailie (HMPPS), Jonathan Van Der Veen (HMPPS), and Tahani Saridar (Think Through Nutrition), whom have all worked towards the betterment of prisoners and their health.

**Financial Support:** This research has in part been supported through the Nigel Groome Scholarship for PhD candidates at Oxford Brookes University

**Conflict of Interest:** The authors acknowledge this work revolves around prison research, where the authors are working in other collaborative projects with HMPPS, and the MOJ, though no conflicts for this review are present.

**Authorship:** This study was conceptualised by M.P, S.C, C.A-M.G, B.L, and J.T, which involved the research question and design. M.P conducted the search strategy and identified papers for screening. Title, abstract, and full text screening was conducted by M.P, S.C, C.A-M.G, and J.T. M.P wrote the initial manuscript, with J.T, S.C, B.L, and C.A-M.G providing comments, and editing of the final draft prior to final manuscript submission.

**Abbreviations:**

ABCL	Adult behaviour checklist
APSD	Antisocial process screening device
AQ	Aggression questionnaire
ASR	Adult self-report
BADDS	Brown's Attention Deficit Disorder Scale
BMI	Body mass index
CI	Confidence interval
CODDS	Conduct and oppositional defiant disorder scales
EPQ	Eysenck personality questionnaire
GHQ-28	General Health Questionnaire-28
HR	Heart rate
HRV	Heart rate variability
IBOS	Inmate behavioural observation scale
IGT	Iowa Gambling Task
IQ	Intelligence quotient
IRM	Institutional reprimand misconduct
IRR	Incident rate ratio
NRCT	Non-randomised controlled trial
POMS	Profile of mood states
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines
PROSPERO	Prospective Register of Systematic Reviews
RCT	Randomised controlled trial
RDA	Recommended daily allowance
rMSSD	Root mean square of successive differences
RoB 2	Revised tool to assess risk of bias in randomised trials
ROBINS-I	Risk of bias in non-randomised studies of interventions

RPQ	Reactive-proactive aggression questionnaire
RR	Rate ratio
SCL-90	Symptom checklist-90
SDAS	Social dysfunction and aggression scale
STAI	State-Trait Anxiety Inventory
ToH	Tower of Hanoi task
WHO	World Health Organisation
YPI	Youth psychopathic trait Inventory

## References

1. World Health Organisation (WHO) (2020) Healthy Diet. [https://www.who.int/news-room/fact-sheets/detail/healthy-diet#:~:text=Consuming%20a%20healthy%20diet%20throughout,diseases%20\(NCDs\)%20and%20conditions](https://www.who.int/news-room/fact-sheets/detail/healthy-diet#:~:text=Consuming%20a%20healthy%20diet%20throughout,diseases%20(NCDs)%20and%20conditions). (accessed 21/10/2023)
2. World Health Organisation (WHO) (2022) Nutrition [https://www.who.int/health-topics/nutrition#tab=tab\\_1](https://www.who.int/health-topics/nutrition#tab=tab_1) (accessed 10/02/2023)
3. World Health Organisation (WHO) (2019) *Essential nutrition actions: mainstreaming nutrition through the life-course*. Geneva: WHO.
4. Geissler C & Powers H (2017) *Human Nutrition*, 13th ed. Oxford: Oxford University Press.
5. World Health Organisation (WHO) (2021) Malnutrition Fact Sheet. [www.who.int/news-room/fact-sheets/detail/malnutrition](http://www.who.int/news-room/fact-sheets/detail/malnutrition) (accessed 11/11/2022)
6. World Health Organisation (WHO) (2021) Obesity and overweight. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> (accessed 21/10/2023)
7. Centers for Disease Control and Prevention (2022) Poor Nutrition. *Chronic Disease Fact Sheets*. <https://www.cdc.gov/chronicdisease/resources/publications/factsheets/nutrition.htm#:~:text=A%20healthy%20diet%20helps%20children,2%20diabetes%2C%20and%20certain%20cancers>. (accessed 21/10/2023)
8. Firth J, Gangwisch JE, Borisini A *et al.* (2020) Food and mood: how do diet and nutrition affect mental wellbeing? *Bmj* 369, m2382.

9. World Health Organisation (WHO) (2021) Mental Health: Key Facts. <https://www.who.int/news-room/fact-sheets/detail/mental-health-strengthening-our-response> (accessed 21/10/2023)
10. World Health Organisation (WHO) (2022) Mental Disorders. <https://www.who.int/news-room/fact-sheets/detail/mental-disorders> (accessed 14/08/2023)
11. NHS (2023) Living with - Depression in adults. (accessed 21/10/2023)
12. The Association of UK Dietitians (2020) Depression and diet. <https://www.bda.uk.com/resource/depression-diet.html> (accessed 21/10/2023)
13. Royal College of Psychiatrists (2020) Depression in adults. <https://www.rcpsych.ac.uk/mental-health/mental-illnesses-and-mental-health-problems/depression#:~:text=Depression%20can%20be%20triggered%20by,breakdown%2C%20or%20losing%20a%20job.&text=If%20your%20life%20circumstances%20mean,more%20likely%20to%20become%20depressed.> (accessed 21/10/2023)
14. Rahe C, Unrath M Berger K Dietary patterns and the risk of depression in adults: a systematic review of observational studies.
15. Li Y, Lv M-R, Wei Y-J *et al.* (2017) Dietary patterns and depression risk: A meta-analysis. *Psychiatry Research* 253.
16. Midboe AM, Gray C, Cheng H *et al.* (2020) Implementation of health-focused interventions in vulnerable populations: protocol for a scoping review. *BMJ Open* 10, e036937.
17. AJMC (2006) Vulnerable Populations: Who Are They? *Supplements and Featured Publications* 12.
18. Fair H & Walmsley R (2021) *World Prison Population List 13th Edition*. World Prison Population List.
19. World Health Organisation (WHO) (2014) *Prisons and Health*. WHO Regional Office for Europe, Denmark: WHO Regional Office for Europe.
20. House of Commons Justice Committee (2021) Mental health in prison: Fifth Report of Session 2021–22, 5th Report ed. [House of Commons, editor]. Westminster: House of Commons.
21. Williams K, Papadopoulou V Booth N (2012) Prisoners' childhood and family backgrounds [Ministry of Justice, editor]. Ministry of Justice: Ministry of Justice.

22. Ministry of Justice (MOJ) (2023) Prison population figures: 2023: Population bulletin: weekly 20th October 2023 [MoJ, editor]. London (UK): Ministry of Justice (MOJ)
23. Goomany A & Dickinson T (2015) The influence of prison climate on the mental health of adult prisoners: a literature review. *J Psychiatr Ment Health Nurs* 22, 413-422.
24. Nurse J, Woodcock P Ormsby J (2003) Influence of environmental factors on mental health within prisons: focus group study. *BMJ* 327, 480.
25. Smoyer AB & Kjær Minke L (2015) *Food systems in correctional settings: a literature review and case study*. WHO Regional Office for Europe, Denmark: World Health Organization. Regional Office for Europ.
26. Stanikowski P, Michalak-Majewska M, Domagała D *et al.* (2020) Implementation of Dietary Reference Intake Standards in Prison Menus in Poland. *Nutrients* 12.
27. Mommaerts K, Lopez NV, Camplain C *et al.* (2022) Nutrition availability for those incarcerated in jail: Implications for mental health. *Int J Prison Health* ahead-of-print.
28. Edwards J, Hartwell H, Reeve W *et al.* (2007) The diet of prisoners in England. *British Food Journal* 109, 216-232.
29. Woods-Brown C, Hunt K Sweeting H (2023) Food and the prison environment: a meta-ethnography of global first-hand experiences of food, meals and eating in custody. *Health Justice* 11, 23.
30. American Psychiatric Association (2022) Warning Signs of Mental Illness. <https://www.psychiatry.org/patients-families/warning-signs-of-mental-illness>. (accessed 21/10/2023)
31. MayoClinic (2023) Self-injury/cutting. <https://www.mayoclinic.org/diseases-conditions/self-injury/symptoms-causes/syc-20350950> (accessed 21/10/2023)
32. NHS (2023) Why People Self-Harm. <https://www.nhs.uk/mental-health/feelings-symptoms-behaviours/behaviours/self-harm/why-people-self-harm/> (accessed 21/10/2023)
33. World Health Organisation (WHO) (2021) Mental health of adolescents. <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health> (accessed 21/10/2023)
34. Gov.uk (2023) Safety and order. *Prisons Data*. <https://data.justice.gov.uk/prisons/safety-and-order> (accessed 21/10/2023)

35. European Prison Observatory (2019) *Prisons in Europe. 2019 Report on European Prisons and Penitentiary Systems*. Online: European Prison Observatory.
36. Office of the Inspector of Custodial Services (2014) *Assaults on staff in Western Australian prisons* [Office of the Inspector of Custodial Services, editor]. Online: Government of Western Australia.
37. Marzano L, Hawton K, Rivlin A *et al.* (2016) Prevention of Suicidal Behavior in Prisons. *Crisis* 37, 323-334.
38. Penal Reform International (2022) *Deaths in prison: Examining causes, responses, and prevention*. Online. <https://www.penalreform.org/resource/deaths-in-prison-examining-causes-responses-and-prevention/>. (Accessed 21/10/2023)
39. US Department of Justice (2019) Press Release: Nearly a Fifth of State and Federal Prisons and a Tenth of Local Jails had at Least One Suicide in 2019 [US Department of Justice, editor]. Online.
40. Page MJ, McKenzie JE, Bossuyt PM *et al.* (2021) The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Bmj* 372, n71.
41. Sterne JAC, Savović J, Page MJ *et al.* (2019) RoB 2: a revised tool for assessing risk of bias in randomised trials. *Bmj* 366, 14898.
42. Sterne JA, Hernán MA, Reeves BC *et al.* (2016) ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 355, i4919.
43. Bachorowski J-A, Newman J, Gans D *et al.* (1994) Effects of sucrose ingestion on juvenile offenders with low, borderline, and normal nadir serum glucose values. *Personality and Individual Differences* 17, 833-843.
44. Cortie CH, Byrne MK, Collier C *et al.* (2020) The Effect of Dietary Supplementation on Aggressive Behaviour in Australian Adult Male Prisoners: A Feasibility and Pilot Study for a Randomised, Double Blind Placebo Controlled Trial. *Nutrients* 12.
45. D'Asaro B, Groesbeck C Nigro C (1975) Diet vitamin program for jail inmates. *Journal of Orthomolecular Psychiatry* 4, 212-222.
46. Gesch CB, Hammond SM, Hampson SE *et al.* (2002) Influence of supplementary vitamins, minerals and essential fatty acids on the antisocial behaviour of young adult prisoners. Randomised, placebo-controlled trial. *Br J Psychiatry* 181, 22-28.
47. Hansen AL, Olson G, Dahl L *et al.* (2014) Reduced anxiety in forensic inpatients after a long-term intervention with Atlantic salmon. *Nutrients* 6, 5405-5418.




48. Hansen AL, Dahl L, Olson G *et al.* (2015) A long-term fatty fish intervention improved executive function in inpatients with antisocial traits and a history of alcohol and drug abuse. *Scand J Psychol* 56, 467-474.
49. Johnson RA, Milner KA, Heng C *et al.* (2018) Implementation and Evaluation of a Physical Activity and Dietary Program in Federal Incarcerated Females. *J Correct Health Care* 24, 395-406.
50. Raine A, Leung C-C, Singh M *et al.* (2020) Omega-3 supplementation in young offenders: A randomized, stratified, double-blind, placebo-controlled, parallel-group trial. *Journal of Experimental Criminology* 16, 389–405.
51. Schoenthaler Stephen Amos Walter Doraz Mary-Ann Kelly George Muedeking James Wakefield Jr S (1997) The effect of randomized vitamin-mineral supplementation on violent and non-violent antisocial behavior among incarcerated juveniles. *Journal of Nutritional & Environmental Medicine* 7, 343-352.
52. Schoenthaler S, Gast D, Giltay EJ *et al.* (2021) The Effects of Vitamin-Mineral Supplements on Serious Rule Violations in Correctional Facilities for Young Adult Male Inmates: A Randomized Controlled Trial. *Crime & Delinquency* 0, 0011128721989073.
53. Zaalberg A, Nijman H, Bulten E *et al.* (2010) Effects of nutritional supplements on aggression, rule-breaking, and psychopathology among young adult prisoners. *Aggress Behav* 36, 117-126.
54. Qureshi FM, Kunaratnam N, Kolla NJ *et al.* (2021) Nutritional supplementation in the treatment of violent and aggressive behavior: A systematic review. *Aggress Behav* 47, 296-309.
55. Meyer BJ, Byrne MK, Collier C *et al.* (2015) Baseline omega-3 index correlates with aggressive and attention deficit disorder behaviours in adult prisoners. *PLoS One* 10, e0120220.
56. Hibbeln JR, Ferguson TA, Blasbalg TL (2006) Omega-3 fatty acid deficiencies in neurodevelopment, aggression and autonomic dysregulation: opportunities for intervention. *Int Rev Psychiatry* 18, 107-118.
57. Micha R, Khatibzadeh S, Shi P *et al.* (2014) Global, regional, and national consumption levels of dietary fats and oils in 1990 and 2010: a systematic analysis including 266 country-specific nutrition surveys. *BMJ : British Medical Journal* 348, g2272.



58. Eves A & Gesch B (2003) Food provision and the nutritional implications of food choices made by young adult males, in a young offenders' institution. *Journal of Human Nutrition and Dietetics* 16, 167-179.
59. Johnson C, Labbé C, Lachance A *et al.* (2022) The Menu Served in Canadian Penitentiaries: A Nutritional Analysis. *Nutrients* 14.
60. Public Health England (PHE) (2016) Government Dietary Recommendations [PHE (PHE), editor]. London (UK): Public Health England (PHE).
61. Wang X, Jiao X, Xu M *et al.* (2020) Effects of circulating vitamin D concentrations on emotion, behavior and attention: A cross-sectional study in preschool children with follow-up behavior experiments in juvenile mice. *Journal of Affective Disorders* 275, 290-298.
62. Robinson SL, Marín C, Oliveros H *et al.* (2020) Vitamin D Deficiency in Middle Childhood Is Related to Behavior Problems in Adolescence. *The Journal of Nutrition* 150, 140-148.
63. Marx W, Manger SH, Blencowe M *et al.* (2023) Clinical guidelines for the use of lifestyle-based mental health care in major depressive disorder: World Federation of Societies for Biological Psychiatry (WFSBP) and Australasian Society of Lifestyle Medicine (ASLM) taskforce . . *The World Journal of Biological Psychiatry* 23, 333-386.
64. Jorde R, Sneve M, Figenschau Y *et al.* (2008) Effects of vitamin D supplementation on symptoms of depression in overweight and obese subjects: randomized double blind trial. *J Intern Med* 264, 599-609.
65. Vellekkatt F & Menon V (2019) Efficacy of vitamin D supplementation in major depression: A meta-analysis of randomized controlled trials. *J Postgrad Med* 65, 74-80.
66. Spedding S (2014) Vitamin D and depression: a systematic review and meta-analysis comparing studies with and without biological flaws. *Nutrients* 6, 1501-1518.
67. Erkkilä AT, Schwab US, de Mello VDF *et al.* Effects of fatty and lean fish intake on blood pressure in subjects with coronary heart disease using multiple medications.
68. Deane KHO, Jimoh OF, Biswas P *et al.* Omega-3 and polyunsaturated fat for prevention of depression and anxiety symptoms: systematic review and meta-analysis of randomised trials.

69. Przybelski RJ & Binkley NC (2007) Is vitamin D important for preserving cognition? A positive correlation of serum 25-hydroxyvitamin D concentration with cognitive function. *Arch Biochem Biophys* 460, 202-205.
70. Wilkins CH, Sheline YI, Roe CM *et al.* (2006) Vitamin D deficiency is associated with low mood and worse cognitive performance in older adults. *Am J Geriatr Psychiatry* 14, 1032-1040.
71. Kruesi MJ, Rapoport JL, Cummings EM *et al.* (1987) Effects of sugar and aspartame on aggression and activity in children. *Am J Psychiatry* 144, 1487-1490.
72. Prinz RJ, Roberts WA, Hantman E (1980) Dietary correlates of hyperactive behavior in children. *J Consult Clin Psychol* 48, 760-769.
73. Wolraich ML, Wilson DB, White JW (1995) The Effect of Sugar on Behavior or Cognition in Children: A Meta-analysis. *JAMA* 274, 1617-1621.
74. Benton D & Stevens MK (2008) The influence of a glucose containing drink on the behavior of children in school. *Biological Psychology* 78, 242-245.
75. Mahoney CR, Taylor HA, Kanarek RB (2007) Effect of an afternoon confectionery snack on cognitive processes critical to learning. *Physiology & Behavior* 90, 344-352.
76. Malay S & Chung KC (2012) The choice of controls for providing validity and evidence in clinical research. *Plast Reconstr Surg* 130, 959-965.
77. Uemura M, Hayashi F, Ishioka K *et al.* Obesity and mental health improvement following nutritional education focusing on gut microbiota composition in Japanese women: a randomised controlled trial.
78. Yu HJ, Li F, Hu YF *et al.* (2020) Improving the Metabolic and Mental Health of Children with Obesity: A School-Based Nutrition Education and Physical Activity Intervention in Wuhan, China. *Nutrients* 12.
79. Foster K, Cuzzillo C, Furness T (2018) Strengthening mental health nurses' resilience through a workplace resilience programme: A qualitative inquiry. *J Psychiatr Ment Health Nurs* 25, 338-348.
80. Abiola T & Udofia O Psychometric assessment of the Wagnild and Young's resilience scale in Kano, Nigeria.
81. European Food Safety Authority (EFSA) (2017) *Overview on Dietary Reference Values for the EU population as derived by the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA)*. Summary of Dietary Reference Values – version 4 (September 2017). EFSA.

82. Fair H & Walmsley R (2022) *World Female Imprisonment List*. World Female Imprisonment List. Institute for crime and Justice Policy Research.
83. Brown RJ, Blum K Trachtenberg MC (1990) Neurodynamics of Relapse Prevention: A Neuronutrient Approach to Outpatient DUI Offenders,. *Journal of Psychoactive Drugs* 22, 173-187.
84. Faintuch J, Garcia Soriano F, Ladeira JP *et al.* (2001) Refeeding procedures after 43 days of total fasting. *Nutrition* 17, 100-104.
85. McKinnon I, Lewis T, Mehta N *et al.* (2018) Vitamin D in patients with intellectual and developmental disability in secure in-patient services in the North of England, UK. *BJPsych Bulletin* 42, 24-29.
86. Abdul Rahman S, Jeffree MS, Kamaludin F *et al.* (2017) Beriberi Outbreak Among Unauthorised Immigrants in a Detention Camp in Malaysia. *Journal of Immigrant and Minority Health* 20 pages, 1294–1297.
87. Shniderman AB & Solberg LB Cosmetic Psychopharmacology for Prisoners: Reducing Crime and Recidivism Through Cognitive Intervention.
88. Najaf Ali Soomro NAC, Safdar Ali Amur, Muhammad Haneef Mugheri, Muzna Paras, Quratulain Khuhro. (2016) Factors associated with delinquent behaviour of inmates at Naara jail Hyderabad, Pakistan. *Rawal Medical Journal* 41, 441-445.
89. Stanikowski P, Michalak-Majewska M, Domagała D *et al.* (2020) Implementation of Dietary Reference Intake Standards in Prison Menus in Poland. *Nutrients* 12, 728.
90. Starzomska M (2006) Ethical challenges in forcible feeding among patients with anorexia nervosa and prisoners. *Archives of Psychiatry and Psychotherapy* 8, 85-96.
91. Virkkunen ME, Horrobin DF, Jenkins DK *et al.* (1987) Plasma phospholipid essential fatty acids and prostaglandins in alcoholic, habitually violent, and impulsive offenders. *Biol Psychiatry* 22, 1087-1096.
92. Virkkunen M, Rissanen A, Naukkarinen H *et al.* (2007) Energy substrate metabolism among habitually violent alcoholic offenders having antisocial personality disorder. *Psychiatry Research* 150, 287-295.
93. Zaalberg A (2019) The effects of nutrients and neurotoxicants on aggressive behavior. *Journal of Criminal Justice* 65.



**Figure 1:** PRISMA 2020 Screening flow chart for the systematic review.

**Table 1:** Included study general and participant characteristics.

Author, Year	Intervention Type	Study Design	Location	Group Studied, No. Participants	Mean Age (Years, SD) (Range)	Study Length	Inclusion/Exclusion Criteria
Bachorowski <i>et al.</i> , (1994) <sup>(43)</sup>	Diet Change	RCT	USA	Juvenile male offenders, (n=48)	Mean not provided, (14-19)	3 Day	Inclusion: Juvenile offenders, between 14-19 years old, with parental permission to volunteer for participation. Exclusion: those with medical records contraindicating participation e.g., diabetes.
Cortie <i>et al.</i> , (2020) <sup>(44)</sup>	Supplements	RCT	Australia	Male adult prisoners, (n=131)	Intervention group: 33.7, 12.6 (19-80). Placebo group: 33.3, 10.3 (18-70)	16 Weeks	Inclusion: Participants who were prisoners and 18+
D'Asaro <i>et al.</i> , (1975) <sup>(45)</sup>	Supplements	NRCT	USA	Inmates, (n=25)	N.D	8 Weeks	Inclusion: Participants were expected to be incarcerated for 2 months minimum, and participation must be approved by the prison physicians and Warden.
Gesch <i>et al.</i> , (2002) <sup>(46)</sup>	Supplements	RCT	UK	Male adult prisoners, (n=231)	N.D	9 Months	Inclusion: 18+ years old.
Hansen <i>et al.</i> , (2014) <sup>(47)</sup>	Diet Change	RCT	USA	Male forensic inpatients, (n=95)	Intervention group: 42.76, 9.72 (21-60). Placebo group: 40.84, 9.14 (21-60)	23 Weeks	Exclusion: IQ<75
Hansen <i>et al.</i> , (2015) <sup>(48)</sup>	Diet Change	RCT	USA	Male forensic inpatients, (n=83)	Intervention group: 42.76, 9.72 (21-60). Placebo group: 40.84, 9.14 (21-60)	23 Weeks	Exclusion: IQ<75
Johnson <i>et al.</i> , (2018) <sup>(49)</sup>	Diet Education	Quasi Experimental Pilot	USA	Female prisoners, (n=29)	42.9, 12.0	12 Weeks	Inclusion: 18+ years old, speaks and reads in English, incarcerated >3 weeks but ≤ 12 months, ≥6 months to serve sentence, BMI ≥18 m/kg <sup>2</sup> , and received health clearance from the nurse practitioner (who was also the principal investigator)

Raine <i>et al.</i> , (2020) <sup>(50)</sup>	Supplements	RCT	Singapore	Young male offenders, (n=145)	Intervention group: 12 Months 19.22, 1.10. Placebo group: 19.29, 1.78. Control group 19.23, 1.31.	Inclusion criteria: 16+ years old, residing at the reformatory training centre at the time of the study. Exclusion criteria: fish allergy, use of omega-3 supplements within the past 3 months, intellectual disability, and the ineligible to enter the reformatory training facility
Schoenthaler <i>et al.</i> , (1997) <sup>(51)</sup>	Supplements	RCT	USA	Incarcerated Juveniles, (n=62)	Intervention group: 3 Months 15.2 Placebo group: 15.2	Inclusion: residents within the medical facility, with parental/guardian consent.
Schoenthaler <i>et al.</i> , (2021) <sup>(52)</sup>	Supplements	RCT	USA	Young male inmates, (n=449)	Low-dose group: 15 Weeks 19.5, 1.4. High-dose group: 19.3, 1.5. Placebo group: 19.4, 1.4.	Inclusion: All prisoners 17-24 were eligible
Zaalberg <i>et al.</i> , (2010) <sup>(53)</sup>	Supplements	RCT	The Netherlands	Male adult offenders, (n=221)	21, 1.5 (18-25)	3 Months No specific inclusion or exclusion criteria are given

RCT, randomised controlled trial; IQ, intelligence quotient; BMI, body mass index; NRCT, non-randomised controlled trial; N.D, no data; SD, standard deviation.

**Table 2:** Included study exposure, control, outcomes, and results summary.

Ref	Exposure(s)	Control/Comparator	Outcome(s)	Outcome Measurement Tool(s)	Key Results
(43)	Sucrose-loaded (78g) breakfast, containing: 636 kcal, 80% carbohydrate, 11% protein, and 9% fat	Comparator no sucrose loaded breakfast, containing: 470kcal, 62% carbohydrate, 21% protein, and 17% fat	(1) Behaviour	(1) 32-Behavioural Checklist (Completed by Experimenters)	There was an interaction between nadir serum glucose group, order of breakfast and subscale of the behaviour checklist, indicating participants with low and normal nadirs were rated as having improved behaviours following the sucrose-loaded meal ( $p < 0.05$ ).
(47)	Farmed Atlantic salmon for dinner (grilled or boiled), portion size 150–300 g, three times per week	No salmon, an alternative meal was provided i.e., chicken, pork, beef, etc.	(1) Anxiety (biological) (2) Anxiety (self-report)	(1) HR, (1) rMSSD, (2) STAI	Regular fatty fish consumption improved HR, HRV baseline to post-intervention for those consuming the fish ( $p < 0.01$ ; $d = 0.45$ ). State-anxiety was also found to be reduced ( $p < 0.03$ ; $d = 0.43$ ), though trait-anxiety was found not to be reduced ( $p > 0.05$ ).
(48)	Farmed Atlantic salmon for dinner (grilled or boiled), portion size 150–300 g, three times per week	No salmon, an alternative meal was provided i.e., chicken, pork, beef, etc.	(1) Cognitive function	(1) IGT, (1) ToH	Regular fatty fish consumption may improve executive functions in forensic inpatients with antisocial traits, and a history of substance abuse ( $p = 0.013$ ; $d = 0.45$ ).
(49)	Health education, and intervention monitoring/support with the nurse practitioner	No control	(1) Resilience	(1) Wagnild and Young's Resilience Scale	Health education slightly improved resilience scores baseline (148.50) to post-intervention (150.50), though not significantly ( $p = 0.437$ ).
(44)	Omega 3 capsules containing fish oil (32% EPA and DHA), and a multivitamin supplement	Placebo, and multivitamin	(1) Rule violations (2) Aggression (self-report)	(1) IBOS, (1) IRM, (2) AQ, (2) BADDSS	Supplements were found to reduce aggression and rule violations, though not significantly ( $p > 0.05$ ). Those found aggressive at baseline had a greater reduction in aggression and rule violations, though not significant ( $p > 0.05$ ).
(45)	Vitamin supplements	Placebo and diet education	(1) Abnormal behaviours	(1) Experiential World Inventory, (1) EPQ (neuroticism dimension), (1) Groesbeck Expanded, (1) Inmate Questionnaire	The effects of the vitamin supplements were found to be not significant for all measures apart from 4 out of 8 dimensions on the Experiential World Inventory, for the dimensions of: sensory, time, ideation, and self, indicating a decrease in feelings of depression. No level of significance was provided.
(46)	Vitamin, mineral, and fatty acid supplements	Placebo	(1) Rule violations	(1) Governor/minor reports	Reduction of 26% (95% CI 8.3-44; $p = 0.03$ ) was seen for rule violations, indicating reduction in antisocial behaviours in the supplement group.

(50) Omega-3 supplement 200ml fruit-based drink, containing; 300mg DHA, 300mg EPA, 180mg alpha-linolenic acid, 60mg DPA, 5µg vitamin D, and antioxidants (unspecified quantities)	200ml fruit-based drinking containing 5µg vitamin D, and antioxidants (unspecified quantities). Control group: no drink.	(1) Antisocial/aggressive behaviours, (2) Rule violations	(1) AQ (self-report), (1) YPI (self-report), (1) RPQ (self-report and officer-report), (1) ASR (self-report), (1) CODDS (self-report and officer-report), (1) APSD (self-report and officer-report), (1) ABCL (officer report), (2) SDAS (officer-report), (2) Institutional infractions	Supplements were found to reduce antisocial behaviour, both short and long term, compared to control. After the intervention period (p=0.006; d=0.56), and 6 months post-intervention (p=0.045; d=0.43), based on the AQ self-reported measure. Officer reported infractions were found not significantly different between groups (p>0.36).
(51) Vitamin-mineral supplements containing 12 vitamins (300% USA RDA) and 11 minerals (100% USA RDA)	Placebo	(1) Rule violation	(1) Rule violation reports	Those taking supplements were seen to experience an 83% decrease in non-violent rule violations, compared to the 49% seen in the placebo group (p=0.008). Violent rule violations fell 80% in the supplement group, and 56% in the placebo group (p=0.044). Total rule violations fell 83% for the supplement group, and 55% in the placebo group (95% CI: 15-41%, p=0.005)
(52) Intervention 1: Low-dose vitamin-mineral supplement containing 100% of the recommended daily allowance, by USA standards, of vitamins and minerals. Intervention group 2: High-dose supplements containing the same amount of vitamins A, D, E, K, Folic acid, Biotin, Copper, Iodine, Iron, and Zinc as the low-dose supplement, but had higher doses of the water-soluble B and C vitamins, and addition of Selenium, Chromium, Manganese, and Molybdenum	Placebo	(1) Rule violation, (2) Mood, (3) Personality traits, (4) Intelligence	(1) Rule violation reports, (2) POMS, (3) EPQ	Supplements covering 100% RDA were found to be able to reduce rule violations, compared with the placebo group (RR=0.61, 95% CI: 41-90, p=0.01). There was no significant result found between the placebo and high-dose group (RR=1.48, 95% CI:0.99-2.23, p=0.06).
(53) Mineral-vitamin supplement, omega-3 fatty acids (EPA and DHA), and omega-6 fatty acid g-linolenic acid	Placebo	(1) Aggression, (2) rule violations, (3) psychopathy	(1) AQ, (2) SDAS, (3) GHQ-28, (3) SCL-90	Supplements were found to reduce rule violations, (IRR=0.60; p=0.017). No effect was seen with aggression or psychopathy (p>0.05).

Heart rate, HR; root mean square of successive differences between normal heartbeats, rMSSD; state trait anxiety index, STAI; heart rate variability, HRV; Iowa gambling test, IGT; tower of Hanoi, ToH; inmate behavioural observation scale, IBOS; institutional reprimand misconduct, IRM; aggression questionnaire, AQ; Brown's attention deficit disorder scale, BADDS; Eysenck personality questionnaire, EPQ; confidence interval, CI; Buss Perry Aggression Questionnaire, AQ; youth psychopathic trait Inventory, YPI; reactive-proactive aggression questionnaire, RPQ; adult self-report, ASR; conduct and oppositional defiant disorder scales, CODDS; antisocial process screening device, APSD; adult behaviour checklist, ABCL; social dysfunction and aggression scale, SDAS; recommended daily allowance, RDA; profile of mood states, POMS; rate ratio, RR; general health questionnaire 28, GHQ-28; symptom checklist-90, SCL-90; incident rate ratio, IRR.